CHROMOSOME NUMBER OF THEVETIA AHOUAI (APOCYNACEAE: RAUVOLFOIDAE: PLUMERIEAE) WITH DISCUSSION ON THE GENERIC BOUNDARIES OF THEVETIA

Justin K. Williams and Julia K. Stutzman¹

Department of Biological Sciences Sam Houston State University Huntsville, Texas 77341-2116, U.S.A.

ABSTRACT

The mitotic chromosome count (2n = 20) for Thevetia ahouai is the first reported chromosome count for Thevetia sect. Ahouai. The count together with a previous count in Thevetia sect. Thevetia (also 2n = 20) provides an additional synapomorphy that further supports the monophyly of Thevetia as traditionally recognized. A discussion on the proposal to recognize Thevetia sect. Thevetia as the genus Cascabela is provided.

Key Words: Thevetia, Apocynaceae, chromosome number, Cascabela, Cerbera, Plumerieae

RESUMEN

El recuento cromosomático mitótico (2n = 20) de Thevetia ahouai es el primero para Thevetia sect. Ahouai. Este recuento junto con otro previo en Thevetia sect. Thevetia (también 2n = 20) aporta una sinapomorfía adicional que apoya la monofilia de Thevetia como se ha reconocido tradicionalmente. Se aporta una discusión de la propuesta de reconocer Thevetia sect. Thevetia como el género Cascabela.

Thevetia L. belongs to the Apocynaceae subfamily Rauvolfioideae tribe Plumerieae and comprises eight species of shrubs occurring from Central Mexico to northern South America (Gensel 1969; Williams 1996; Allorge 1998; Endress et al. 2007). According to various specialists in the Apocynaceae, the generic boundaries of Thevetia vary. In the concept of Thevetia sensu K. Schum. (Schumann 1895; Gensel 1969; Williams 1996; Allorge 1998; Alvarado-Cardenas 2004), eight species are sub-divided between two sections: sect. Ahouai K. Schum. with three species—T. ahouai (L.) A. DC., T. amazonica Ducke, and T. bicornuta Mull. Arg.—and sect. Thevetia K. Schum. with five species—T. gaumeri Hemsl., T. ovata (Cav.) A. DC., T. peruviana (Pers.) K. Schum., T. pinifolia (Standl. & Steyerm.) J.K. Williams, and T. thevetiodes (H.B.K.) K. Schum. In the concept of Thevetia sensu Lippold (Lippold 1980; Alvarado-Cardenas & Ochoterena 2007) the three species of sect. Ahouai are retained in Thevetia; the other five species of sect. Thevetia are segregated into the genus Cascablea Raf. Although Lippold (1980) and Alvarado-Cardenas and Ochoterena (2007) segregate Thevetia sensu K. Schum. into the two genera Thevetia and Cascablea, at no point do they argue against the genera's "close morphological relationship" (Alvarado-Cardenas & Ochoterena 2007). In fact, a recent morphological cladistic analysis (Fig. 1, Alvarado-Cardenas & Ochoterena 2007) nests all eight species of Thevetia sensu K. Schum. in a clade supported by six synapomorphies. In short, Thevetia sensu K. Schum. is clearly shown to be monophyletic. Nevertheless, Alvarado-Cardenas and Ochoterena (2007) argue for its paraphyly and choose instead to recognize the two sub-clades of the clade (Fig. 1) as distinct genera: Thevetia and Cascabela.

Chromosome numbers have proven useful in resolving generic relationships in the Apocynaceae (Van der Laan & Arends 1985; Williams 2007). To date, 73 of the 179 genera of the Apocynaceae s.str. (subfamilies Rauvolfioideae and Apocynoidoideae) have been counted (Van der Laan & Arends 1985; Goldblatt & Johnson 2003; Williams 2007). Although previous chromosome counts for sect. *Thevetia* (T. peruviana; T0 = 20) exist, none have been reported for sect. *Ahouai*. The present paper provides the first reported chromosome count for a species of sect. *Ahouai* and discusses the utility of chromosome numbers in interpreting the systematic relationship of *Thevetia* and *Cascabela*.

¹Current Address: Bridgewater College, Box 1568, Bridgewater, Virginia 22812, U.S.A.




Fig. 1. Morphologically constructed dendogram of "Plumerieae" clade (from Alvarado-Cardenas & Ochoterena 2007). a = Thevetia sensu K. Schum. and/or "Cascabela-Thevetia" clade of Alvarado-Cardenas and Ochoterena 2007. This clade is supported by seven synapomorphies. b = Thevetia sect. Ahouai and/or Thevetia sensu Lippold. This clade is supported by two synapomorphies. c = Thevetia sect. Thevetia and/or Cascablea Lippold. This clade is supported by four synapomorphies.

MATERIALS AND METHODS

Roots tips and voucher specimens were collected from a greenhouse specimen of *Thevetia ahouai* housed at the greenhouse of the Department of Biological Sciences, Sam Houston State University (Table 1). The root tips were fixed and analyzed for chromosome number using standard procedures (Raffauf 1964; Van der Laan & Arends 1985). A voucher specimen of the greenhouse plant was made and is preserved in the Warner Herbarium (SHST).

RESULTS

The format used for reporting chromosome numbers in this article follows that established by Strother and Nesom (1997). A mitotic chromosome number of 2n = 20 was recorded for *Thevetia ahouai*. Van der Laan and Arends (1985) reported chromosome lengths in the Apocynaceae to be between 0.5–4.0 µm, with the average chromosome length between 1–2 µm. The length of the chromosomes in *T. ahouai* varied between 1–3 µm, consistent with most other chromosomes in the Apocynaceae. The base chromosome number (x = 10) for *Thevetia ahouai* is consistent with previous reports of x = 10 for *Thevetia* (x = 10) for *Thevetia* (x = 10). Gadella 1977; Ugborogho 1983; Van der Laan & Arends 1985; Santhosh & Omanakumari 1997).

DISCUSSION

Van der Laan and Arends (1985) postulated a base chromosome number of x = 11 for the Apocynaceae s. str. based on its prevalence in the family and on the observation that many of the plesiomorphic taxa possess a base number of x = 11. A base chromosome number of x = 10 is found in four genera representing three of the 11 tribes recognized in the Rauvolfioideae (Endress et al. 2007): Hunterieae (*Gonioma* E. Mey.), Plumerieae (*Cerbera* L., *Thevetia*), and Vinceae (*Ochrosia* Juss.). Based on tribal circumscription (Endress et al. 2007) along with molecular evidence (Simóes et al. 2007), x = 10 is reconstructed as having evolved independently at least three times in the Rauvolfioideae.

A cursory review of chromosome counts for the Apocynaceae (Van der Laan & Arends 1985; Goldblatt & Johnson 2003) reveals that at present the only chromosome counts for genera in the Plumerieae, and thus relatives to Thevetia (2n = 20), are for Allamanda L. (n = 9; 2n = 18), Cerbera (2n = 40), Himatanthus Willd. ex Schult. (2n = 18), Mortoniella Woodson (2n = 32), and Plumeria L. (2n = 36). At present there are no chromosome counts for the remaining four genera in the Plumerieae: Anechites Griseb., Cameraria L., Cerberiopsis Vieill. ex Pancher & Sébert, and Shytanthus Meyen., and it is suggested here that effort be made to obtain count of these taxa. Figures 1 and 2 present cladograms of the Plumerieae constructed from morphological (Alvarado-Cardenas & Ochoterena 2007) and molecular evidence (Simóes et al. 2007), respectively. Diploid counts for the respective genera included in the phylogenies are presented for both Figures 1 and 2.

Alvarado-Cardenas and Ochoterena (2007) presented six synapomorphies that described the *Thevetia* sensu K. Schum. clade (Fig. 1, branch a). The diploid count of 2n = 20 presented here adds a seventh synapomorphy. When interpreting their data Alvarado-Cardenas and Ochoterena (2007) state that "(t)here is still no consensus regarding the question of whether one should recognize one genus with two (sections) or two distinct genera (Lippold 1980), given that *Cascabela* and *Thevetia* are sister taxa." We would argue that the <u>consensus</u> in evolutionary systematics is to assign generic boundaries that reflect both monophyly and shared ancestry. The decision to divide a well supported clade into two separate genera may support monophyly; however, it excludes shared ancestry. Without prior knowledge, most botanists would be unaware that *Thevetia* and *Cascabela* are sister taxa that share seven synapomorphies. Instead, an evolutionarily more meaningful interpretation of the clade would be that *Thevetia* sensu K. Schum. is monophyletic and support is provided for the recognition of two sections as defined by Schumann (1895). In order to maintain systematic consistency as pertains to current trends in phylogenetic nomenclature we recognize *Thevetia* sensu K. Schum. and regard *Cascabela* and all taxa pertaining to the genus as synonyms of *Thevetia*.

Taxon	Voucher specimen	Chromosome number (2n)	
Thevetia ahouai	TEXAS: Sam Houston State University Department of Biological Sciences Greenhouse specimen, 22 Jan 2008 Williams 2008-1 (SHST).	20	

TABLE 1. Voucher specimen for the chromosome number of Thevetia ahouai.




Fig. 2. Molecularly constructed dendogram of "Plumerieae" clade (from Simóes 2007).

ACKNOWLEDGMENTS

We thank Tami Cook for providing us with access to her digital light microscope, and Bob Rhodes for mixing the Carnoy's solution and aceto-orcein. Sibyl Buceli and an anonymous reviewer provided valuable editorial comments.

REFERENCES

Allorge, L. 1998. Les *Thevetia*, compagnons des succulentes. Succulentes 21(1):23–32.

Alvarado-Cardenas, L.O. 2004. Apocynaceae. Flora del Valle de Tehuacan-Cuicatlan 38:1–57.

ALVARADO-CARDENAS, L.O. and H. Ochoterena. 2007. A phylogenetic analysis of the *Cascabela-Thevetia* species complex (Plumerieae, Apocynaceae) based on morphology. Ann. Missouri Bot. Gard. 94:298–323. 2007.

Endress, M.E. and P. Bruyns. 2000. A revised classification of the Apocynaceae s.l. Bot. Rev. 66:1–56.

Endress, M.E., S. Liede-Schumann, and U. Meve. 2007. Advances in Apocynaceae: The enlightenment, an introduction. Ann. Missouri Bot. Gard. 94:259–267.

Gadella, T.W.J. 1977. IOPB chromosome number reports LVI. Taxon 26:257–274.

Gensel, W.H. 1969. A revision of the genus *Thevetia* (Apocynaceae). Masters Thesis University of Connecticut.

Goldblatt, P. and D.E. Johnson. 2003. Index to plant chromosome numbers 1998–2000. Monogr. Syst. Bot. Missouri Bot. Gard. 94.

Lippold, H. 1980. Die Gattungen *Thevetia L., Cerbera L.* und *Cascabela* Raf. (Apocynaceae). Feddes Repert. 91:45–55.

RAFFAUF, R.F. 1964. Some chemotaxonomic considerations in the Apocynaceae. Lloydia 27:288–298.

Santhosh, B. and N. Omanakumari. 1997. Karyomorphological studies on two varieties of *Thevetia peruviana*. J. Cytol. Gene. 32:95–98.

Schumann, K. 1895. Apocynaceae. In: A. Engler and K.A. Prantl, Die Natürlichen Pflanzenfamilien. Wilhelm Engelmann, Leipzig. 4(2):109–189.

SIMÓES, A.O., T. LIVSHULTZ, E. CONTI, and M.E. ENDRESS. 2007. Phylogeny and systematic of the Rauvolfioideae (Apocynaceae) based on molecular and morphological evidence. Ann. Missouri Bot. Gard. 94:268–297.

Strother, J.L. and G.L. Nesom. 1997 Conventions for reporting plant chromosome numbers. Sida 17:829–831. Ugborogho, R.E. 1983. IOPB chromosome number reports LXXIX. Taxon 32:321.

Van der Laan, F.N. and J.C. Arends. 1985. Cytotaxonomy of the Apocynaceae. Genetica 68:3–35.

Williams, J.K. 1996. A new combination in *Thevetia* (Apocynaceae). Sida 17:185–190.

Williams, J.K. 2007. Documented chromosome numbers 2007: Chromosome number of *Laubertia contorta* (Apocynaceae: Apocynoideae) and its phylogenetic importance. J. Bot. Res. Inst. Texas 1:431–435. 2007.